



Production advice ware potatoes TRADITIONAL FRESH

- Early maturity and tuberisation
- Big size tubers
- Broad adaptation
- Easy to grow
- Good dry matter content



Agronomic characters

Maturity	78	Early
Dormancy	59	Medium
Yield early	94	Good
Yield mature	94	Good
Tuber size	86	Very large
Tuber shape		Round oval / Oval
Number of tubers		6-8
Skin colour		Yellow
Flesh after cooking		Light yellow
Cooking type		AB - Slightly firm
Dry matter content/Starch		19,2% / 13,4%
UWW/Specific gravity		350 / 1,074
Internal bruising	4	Not sensitive
Metribuzin sensitivity	40	● ● ● ● ● ●
Little Potato disorder	52	● ● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	51.000	26	22
35/45	48.000	28	23
35/55	46.000	29	24
50/55	43.000	31	26
55/65	41.500	32	27

Resistances

Foliage Blight	49	● ● ● ● ● ●
Tuber Blight	75	● ● ● ● ● ●
Alternaria	56	● ● ● ● ● ●
Common scab	58	● ● ● ● ● ●
Powdery scab	53	● ● ● ● ● ●
Spraing	84	● ● ● ● ● ●
PVY	21	● ● ● ● ● ●
Yntn tuber tolerance	89	● ● ● ● ● ●

PCN Resistance

Ro1/4 *2	● ● ● ● ● ●
Ro2/3 *1	● ● ● ● ● ●
Ro5 *1	● ● ● ● ● ●
Pa2 *3	● ● ● ● ● ●
Pa3 *2	● ● ● ● ● ●

Wart disease

F1	10	● ● ● ● ● ●
F2		● ● ● ● ● ●
F6		● ● ● ● ● ●
F18		● ● ● ● ● ●

Fertilizer

- Adapt fertilization to soil analysis.
- Always refer to the current local rules about crop fertilization.
- Nitrogen (N) : Very high inputs (>250Kg N/ha inclusive of soil supply)
- A good phosphate and potassium fertilization can increase the yield.

* HZPC own analysis/no official analysis



Production advice ware potatoes TRADITIONAL FRESH



Pre-treatment and planting

- The dormancy is medium long, a heat treatment is not necessary.
- De-sprouting will increase the risk of little potato disorder.
- Do not plant in cold soils as this will increase the risk of irregular emergence.




Growing attention points

- The tuberisation is early.
- Tuber number is quite low and tubers easily get too big.
- JAERLA is susceptible to metribuzin, also when applied before emergence.
- Use a robust Phytophthora programme, to prevent foliage blight.
- Preventive spraying against Phytophthora is advised.



Haulm killing and harvest

- JAERLA is quite early maturing, haulm killing shouldn't give any problems.
- In general, JAERLA has a good stolone detachment and early skin set.



Storage

- For storage period longer than 3 months, refrigeration is required.
- Ventilate regularly, but briefly, to prevent CO2 accumulation.
- Good to storable at constant temperature.