



Production advice ware potatoes France

- Uniform tuber size and shape
- Good bruising tolerance
- Good cooking quality
- Good resistance to common scab
- Good storability



Agronomic characters

Maturity	56	Medium late
Dormancy	60	Medium
Yield early	101	Moderate high
Yield mature	100	Moderate high
Tuber size	81	Large
Tuber shape		Oval
Number of tubers		9-11
Skin colour		Red
Flesh after cooking		Yellow
Cooking type		AB - Slightly firm
Dry matter content/Starch	20,3% / 14,4%	
UWW/Specific gravity	371 / 1,079	
Internal bruising	5	Not sensitive
Metribuzin sensitivity	76	● ● ● ● ●
Little Potato disorder	83	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Market: 45-75 mm.

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	45.000	30	25
35/45	38.000	35	29
45/50	34.000	39	33
50/55	30.000	44	37

Plant depth: normal.

Always check the tuber count for an accurate calculation.

Resistances

Foliage Blight	64	● ● ● ● ●
Tuber Blight	74	● ● ● ● ●
Alternaria	69	● ● ● ● ●
Common scab	72	● ● ● ● ●
Powdery scab	69	● ● ● ● ●
Spraing	98	● ● ● ● ●
PVY	69	● ● ● ● ●
Yntn tuber tolerance	99	● ● ● ● ●

PCN Resistance

Ro1/4	9	● ● ● ● ●
Ro2/3	*1	● ● ● ● ●
Ro5	*1	● ● ● ● ●
Pa2		● ● ● ● ●
Pa3		● ● ● ● ●

Wart disease

F1	10	● ● ● ● ●
F2	3	● ● ● ● ●
F6		● ● ● ● ●
F18		● ● ● ● ●

* HZPC own analysis/no official analysis

Fertilizer

- Adapt fertilization to soil analysis.
- Late availability of nitrogen will delay senescence and results in poor cooking quality.
- Nitrogen (N) : Medium inputs (170-210Kg N/Ha inclusive of soil supply).



Production advice ware potatoes France



Pre-treatment and planting

- Warm up correctly the seed and plant it with small white sprouts.
- Always treat MOZART seeds against Rhizoctonia.
- MOZART can be planted early but avoid planting in cold conditions.




Growing attention points

- The use of metribuzin is recommended before emergence. When applying post emergence, use the low dose system.
- Preventive spraying against Phytophthora is advised.



Haulm killing and harvest

- Aim for a dry matter content of 19,5 %.
- Only a mature crop will give a good storable product.
- MOZART can be harvested after 120 days.
- MOZART has a thin skin, pay extra attention to skin set.
- Minimum 4 weeks between haulm killing and crop lifting for right skin set.



Storage

- Storage temperature: 6 °C.
- Any fluctuation in temperature, combined with condensation, can result in early sprouting and a high risk of Silver scurf.