



Production advice ware potatoes TRADITIONAL FRESH

- Uniform tuber size and shape
- Good bruising tolerance
- Good cooking quality
- Good resistance to common scab
- Good storability



Agronomic characters

Maturity	56	Medium late
Dormancy	60	Medium
Yield early	101	Moderate high
Yield mature	100	Moderate high
Tuber size	81	Large
Tuber shape		Oval
Number of tubers		9-11
Skin colour		Red
Flesh after cooking		Yellow
Cooking type		AB - Slightly firm
Dry matter content/Starch		20,3% / 14,4%
UWW/Specific gravity		371 / 1,079
Internal bruising	5	Not sensitive
Metribuzin sensitivity	76	●●●●●○
Little Potato disorder	83	●●●●●●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	53.000	25	21
35/45	50.000	27	22
45/50	42.000	32	26
45/55	40.000	33	28
35/55	43.000	31	26
50/55	37.000	36	30
55/65	34.000	39	33

Plant depth: normal.

Always check the tuber count for an accurate


Resistances

Foliage Blight	64	●●●●●○
Tuber Blight	74	●●●●●○
Alternaria	69	●●●●●○
Common scab	72	●●●●●●
Powdery scab	69	●●●●●○
Spraing	98	●●●●●●
PVY	69	●●●●●○
Yntn tuber tolerance	99	●●●●●●
PCN Resistance		
Ro1/4	9	●●●●●●
Ro2/3	*1	●●●●●○
Ro5	*1	●●●●●○
Pa2		●●●●●○
Pa3		●●●●●○
Wart disease		
F1	10	●●●●●●
F2	3	●●●●●○
F6		●●●●●○
F18		●●●●●○

* HZPC own analysis/no official analysis



Production advice ware potatoes TRADITIONAL FRESH




Pre-treatment and planting

- Warm up correctly the seed and plant it with small white sprouts.
- Always treat MOZART seeds against Rhizoctonia.
- MOZART can be planted early but avoid planting in cold conditions.




Growing attention points

- The use of metribuzin is recommended before emergence. When applying post emergence, use the low dose system.
- Preventive spraying against Phytophthora is advised.



Haulm killing and harvest

- Aim for a dry matter content of 19,5 %.
- Only a mature crop will give a good storable product.
- MOZART can be harvested after 120 days.
- MOZART has a thin skin, pay extra attention to skin set.
- Minimum 4 weeks between haulm killing and crop lifting for right skin set.



Storage

- Storage temperature: 6 °C.
- Any fluctuation in temperature, combined with condensation, can result in early sprouting and a high risk of Silver scurf.

Plant populations

calculation.

Fertilizer

- Adapt fertilization to soil analysis.
- Late availability of nitrogen will delay senescence and results in poor cooking quality.
- Nitrogen (N) : Medium inputs (170-210Kg N/Ha inclusive of soil supply).

