



General production advice ware potatoes

- Firm cooking
- Uniform tuber size and shape
- Good tolerance to dry circumstances
- Good bruising tolerance
- Good storability



Agronomic characters

Maturity	61	Medium late
Dormancy	91	Very long
Yield mature	83	Moderate
Tuber size	66	Small
Tuber shape		Long oval / Long
Number of tubers		12-14
Skin colour		Yellow
Flesh after cooking		Yellow
Cooking type		A - Firm
Dry matter content/Starch		19,4% / 14,2%
UWW/Specific gravity		367 / 1,078
Internal bruising	4	Not sensitive
Metribuzin sensitivity	78	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Market: 35-55 mm.

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
30/40	44.000	30	25
40/45	42.000	32	26
45/50	40.000	33	28

Resistances

Foliage Blight	52	● ● ● ● ●
Tuber Blight	70	● ● ● ● ●
Common scab	45	● ● ● ● ●
Powdery scab	58	● ● ● ● ●
Spraing	99	● ● ● ● ●
PVY	70	● ● ● ● ●
Yntn tuber tolerance	99	● ● ● ● ●

PCN Resistance

Ro1/4 *1	● ● ● ● ●
Ro2/3	● ● ● ● ●
Ro5	● ● ● ● ●
Pa2 *1	● ● ● ● ●
Pa3 *1	● ● ● ● ●

Wart disease

F1	● ● ● ● ●
F2	● ● ● ● ●
F6	● ● ● ● ●
F18	● ● ● ● ●

* HZPC own analysis/no official analysis

Fertilizer

- Always refer to the local and current rules about crop fertilization.
- Nitrogen (N) : Medium inputs (170-210Kg N/ha inclusive of soil supply).



General production advice ware potatoes



Pre-treatment and planting

- NORMANDELINE has a good dormancy.
- Warm up correctly the seed and plant it with small white sprouts.
- Always treat NORMANDELINE seeds against Rhizoctonia.
- Plant into warm soils, don't plant too early.
- Avoid soils with high common scab risk.




Growing attention points

- Emergence is somewhat slow, later on foliage develops well.
- NORMANDELINE has a good tolerance to metribuzin.
- Preventive spraying against Phytophthora is advised.



Haulm killing and harvest

- Minimum 3 weeks between haulm killing and crop lifting for right skin set.
- Prevent mechanical damage to improve storability.



Storage

- With a proper store management, NORMANDELINE is suitable for long term storage.
- Storage temperature: 4 °C.
- As NORMANDELINE has a long dormancy, the quantity of sprout inhibitors can be reduced for long term storage.