



Production advice ware potatoes TRADITIONAL FRESH

- Very attractive bright skin
- Good yield
- Good dry matter content
- Good resistance to common scab



Agronomic characters

Maturity	61	Medium late
Dormancy	96	Very long
Yield mature	115	Very high
Tuber size	82	Large
Tuber shape		Oval
Number of tubers		9-11
Skin colour		Yellow
Flesh after cooking		Yellow
Cooking type		AB - Slightly firm
Dry matter content/Starch		19% / 13,2%
UWW/Specific gravity		346 / 1,074
Internal bruising	15	Quite sensitive
Metribuzin sensitivity	67	● ● ● ● ●
Little Potato disorder	90	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	53.000	25	21
35/45	48.000	28	23
35/55	44.000	30	25
50/55	40.000	33	28
55/65	38.000	35	29

Plant depth: normal.

Always check the tuber count for an accurate calculation.

Resistances

Foliage Blight	42	● ● ● ● ●
Tuber Blight	60	● ● ● ● ●
Common scab	61	● ● ● ● ●
Powdery scab	59	● ● ● ● ●
Spraing	97	● ● ● ● ●
PVY	18	● ● ● ● ●
Yntn tuber tolerance	99	● ● ● ● ●

PCN Resistance

Ro1/4	9	● ● ● ● ●
Ro2/3	*1	● ● ● ● ●
Ro5	*8	● ● ● ● ●
Pa2		● ● ● ● ●
Pa3		● ● ● ● ●

Wart disease

F1	1	● ● ● ● ●
F2	*1	● ● ● ● ●
F6	*1	● ● ● ● ●
F18	*1	● ● ● ● ●

* HZPC own analysis/no official analysis

Fertilizer

- Adapt fertilization to soil analysis.
- Always refer to the local and current rules about crop fertilization.
- Apply 2/3 before planting and 1/3 as top dressing.
- Nitrogen (N) : Medium inputs (170-210Kg N/Ha inclusive of soil supply).
- Phosphate (P) and Potassium (K) : according to standard advice.
- Pay attention to the amount of organic fertilizer to avoid later mineralisation.



Production advice ware potatoes **TRADITIONAL FRESH**



Pre-treatment and planting

- PERDIZ has a long dormancy.
- Pre-sprouting helps to advance the growth.
- PERDIZ can be grown on all soil types.
- Can be planted with small white buds but good pre-sprouting can advance the crop.




Growing attention points

- Emergence is uniform with a good foliage development.
- In spite of the late maturity, PERDIZ has an early tuberisation.
- PERDIZ is quite sensitive to metribuzin, only use it pre-emergence.
- Use a robust Phytophthora programme, to prevent foliage and tuber blight



Haulm killing and harvest

- Ensure a sufficient matured crop before haulm killing. This promotes a higher dry matter content.
- Tubers are big size. Harvest and store with care to prevent damage and bruising.
- The skin should be fully set before harvest.
- Prevent mechanical damage to improve storability.



Storage

- PERDIZ has a long dormancy and has good storability.
- Dry quickly after harvest and keep dry to help prevent silver scurf.
- Stable temperature during storage prevents silver scurf development.
- As PERDIZ has a long dormancy, the quantity of sprout inhibitors can be reduced for long term storage.