




Production advice ware potatoes TRADITIONAL FRESH

- Very attractive bright skin
- High yield
- Good dry matter content
- Suitable to make (home) fries
- Good common and powdery scab resistance



Agronomic characters

Maturity	57	Medium late
Dormancy	66	Medium
Yield mature	119	Very high
Tuber size	86	Very large
Tuber shape		Oval
Number of tubers		9-11
Skin colour		Yellow
Flesh after cooking		Yellow
Cooking type		AB - Slightly firm
Dry matter content/Starch		19,1% / 13,3%
UWW/Specific gravity		348 / 1,074
Internal bruising	16	Sensitive
Metribuzin sensitivity	73	● ● ● ● ●
Little Potato disorder	75	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	50.000	27	22
35/45	44.000	30	25
35/55	42.000	32	26
50/55	39.000	34	28
55/65	36.000	37	31

Plant depth: normal.

Always check the tuber count for an accurate calculation.

Resistances

Foliage Blight	53	● ● ● ● ●
Tuber Blight	62	● ● ● ● ●
Alternaria	70	● ● ● ● ●
Common scab	73	● ● ● ● ●
Powdery scab	66	● ● ● ● ●
Spraing	91	● ● ● ● ●
PVY	17	● ● ● ● ●
Yntn tuber tolerance	99	● ● ● ● ●

PCN Resistance

Ro1/4	9	● ● ● ● ●
Ro2/3	3	● ● ● ● ●
Ro5	*1	● ● ● ● ●
Pa2		● ● ● ● ●
Pa3		● ● ● ● ●

Wart disease

F1	*3	● ● ● ● ●
F2	*5	● ● ● ● ●
F6	*5	● ● ● ● ●
F18	*1	● ● ● ● ●

* HZPC own analysis/no official analysis

Fertilizer

- Adapt fertilization to soil analysis.
- Always refer to the local and current rules about crop fertilization.
- Apply 2/3 before planting and 1/3 as top dressing.
- Nitrogen (N) : Medium inputs (170-210Kg N/Ha inclusive of soil supply).
- Phosphate (P) and Potassium (K) : according to standard advice.



Production advice ware potatoes TRADITIONAL FRESH



Pre-treatment and planting

- RASHIDA has a medium dormancy.
- Allow the seed to acclimatize to the local conditions before planting.
- RASHIDA can be grown on all soil types.
- Take care of a good volume ridge to prevent green tubers.




Growing attention points

- Emergence is uniform with a good foliage development.
- RASHIDA has a good tolerance to metribuzin.
- Use a robust Phytophthora programme, to prevent foliage blight.



Haulm killing and harvest

- Ensure a sufficient matured crop before haulm killing. This promotes a higher dry matter content.
- An unset skin will have a negative effect on storability.
- RASHIDA is sensitive to bruising and mechanical damage.



Storage

- RASHIDA has a good dormancy, therefore ambient storage can be used for the first months after harvest.
- Good to storable at constant temperature.