




## General production advice ware potatoes

- High marketable yield
- Uniform tuber size and shape
- Length specifications
- Suitable for long term storage



## Agronomic characters

Maturity	65	Medium early
Dormancy	73	Long
Yield mature	103	Moderate high
Tuber size	79	Medium
Tuber shape		Long oval
Number of tubers		12-14
Skin colour		Yellow
Flesh after cooking		Light yellow
Cooking type		B - Slightly mealy
Dry matter content/Starch		23% / 17%
UWW/Specific gravity		426 / 1,092
Internal bruising	14	Quite sensitive
Metribuzin sensitivity	65	● ● ● ● ●
Little Potato disorder	88	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

## Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	45.000	30	25
35/45	38.000	35	29
35/50	36.000	37	31
45/50	34.000	39	33
50/60	32.000	42	35
50/60 cut seed	38.000	35	29

Plant depth for medium & big seed sizes : 3 cm deeper than normal.

## Resistances

Foliage Blight	50	● ● ● ● ●
Tuber Blight	70	● ● ● ● ●
Alternaria	71	● ● ● ● ●
Common scab	62	● ● ● ● ●
Powdery scab	62	● ● ● ● ●
Spraing	99	● ● ● ● ●
PVY	59	● ● ● ● ●
Yntn tuber tolerance	99	● ● ● ● ●

### PCN Resistance

Ro1/4	9	● ● ● ● ●
Ro2/3	*6	● ● ● ● ●
Ro5	*5	● ● ● ● ●
Pa2		● ● ● ● ●
Pa3		● ● ● ● ●

### Wart disease

F1	8	● ● ● ● ●
F2		● ● ● ● ●
F6		● ● ● ● ●
F18		● ● ● ● ●

\* HZPC own analysis/no official analysis

## Fertilizer

- Nitrogen (N) : 210-250 kg N/ha inclusive of soil supply.
- Apply 2/3 before planting and 1/3 as top dressing.
- Phosphate (P) : according to standard advice.
- Potassium (K): approx 350 kg K/ha



## General production advice ware potatoes




### Pre-treatment and planting

- TRAVOLTA has a long dormancy.
- TRAVOLTA is sensitive to chemical treatment against Rhizoctonia. Do not use chemical treatment against Rhizoctonia on tubers directly before planting and/or when sprouts are visible.
- Take care of a good volume ridge to prevent green tubers.




### Growing attention points

- TRAVOLTA is susceptible to Alternaria. Start treatments at flowering time.
- Preventive spraying against Phytophthora is advised.



### Haulm killing and harvest

- TRAVOLTA is moderately susceptible to bruising. Reduce drop heights and avoid mechanical damage.



### Storage

- Take care of the wound healing period.
- Store at a stable 7 °C.
- For long term storage it is necessary to use sprout inhibitors.
- With a proper store management, TRAVOLTA is suitable for long term storage.