




Production advice ware potatoes France

- Dark yellow flesh
- Good cooking quality
- Multi-purpose variety
- Excellent frying quality
- Good storability



Agronomic characters

Maturity	63	Medium late
Dormancy	70	Long
Yield mature	97	Good
Tuber size	83	Large
Tuber shape		Oval / Long oval
Number of tubers		9-11
Skin colour		Yellow
Flesh after cooking		Yellow
Cooking type		B - Slightly mealy
Dry matter content/Starch		20,9% / 15,1%
UWW/Specific gravity		385 / 1,083
Internal bruising	18	Sensitive
Metribuzin sensitivity	61	● ● ● ● ●
Little Potato disorder	75	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Market: 40-70 mm.

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	40.000	33	28
35/45	36.000	37	31
45/50	30.000	44	37

Plant depth: normal.

Always check the tuber count for an accurate calculation.

Resistances

Foliage Blight	39	● ● ● ● ●
Tuber Blight	70	● ● ● ● ●
Alternaria	76	● ● ● ● ●
Common scab	52	● ● ● ● ●
Powdery scab	47	● ● ● ● ●
Spraing	76	● ● ● ● ●
PVY	39	● ● ● ● ●
Yntn tuber tolerance	96	● ● ● ● ●

PCN Resistance

Ro1/4	8	● ● ● ● ●
Ro2/3	*1	● ● ● ● ●
Ro5	*1	● ● ● ● ●
Pa2	*1	● ● ● ● ●
Pa3	*1	● ● ● ● ●

Wart disease

F1	10	● ● ● ● ●
F2	6	● ● ● ● ●
F6	6	● ● ● ● ●
F18		● ● ● ● ●

* HZPC own analysis/no official analysis

Fertilizer

- Always refer to the local and current rules about crop fertilization.
- Nitrogen (N) : Medium inputs (170-210Kg N/ha inclusive of soil supply).
- Phosphate (P) and Potassium (K) : according to standard advice.



Production advice ware potatoes France



Pre-treatment and planting

- VICTORIA has a good dormancy.
- Allow the seed to acclimatize to the local conditions before planting.
- Always treat VICTORIA seeds against Rhizoctonia.
- Planting with small white sprouts produces the best results.
- Take care of a good volume ridge to prevent green tubers.




Growing attention points

- Emergence is uniform with a good foliage development.
- The tuberisation is late.
- VICTORIA grows spread in the ridge.
- The use of metribuzin is recommended before emergence. When applying post emergence, use the low dose system.
- Weather conditions have a strong influence on metribuzin effectivity, adapt dose to the circumstances.
- Use a robust Phytophthora programme, to prevent foliage blight.



Haulm killing and harvest

- Aim for a dry matter content of 19,5 %.
- Minimum 3 weeks between haulm killing and crop lifting for right skin set.
- VICTORIA is moderately susceptible to bruising. Reduce drop heights and avoid mechanical damage.



Storage

- VICTORIA has a long dormancy and has good storability.
- VICTORIA is suitable for long term storage.
- Dry quickly after harvest and keep dry to help prevent silver scurf.
- REFRIGERATED STORAGE
- Reduce temperature 0,5 - 0,7 degrees a day to a stable value of 6°C.
- For long term storage it is necessary to use sprout inhibitors.