




Production advice ware potatoes FRENCH FRIES

- Dark yellow flesh
- High yield
- Good cooking quality
- Multi-purpose variety
- Consistent high frying quality



Agronomic characters

Maturity	63	Medium late
Dormancy	70	Long
Yield mature	97	Good
Tuber size	83	Large
Tuber shape		Oval / Long oval
Number of tubers		9-11
Skin colour		Yellow
Flesh after cooking		Yellow
Cooking type		B - Slightly mealy
Dry matter content/Starch		20,9% / 15,1%
UWW/Specific gravity		385 / 1,083
Internal bruising	18	Sensitive
Metribuzin sensitivity	61	● ● ● ● ●
Little Potato disorder	75	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	50.000	27	22
35/50	38.000	35	29
50/60	32.000	42	35
50/60 cut seed	40.000	33	28

Plant depth: normal.

Resistances

Foliage Blight	39	● ● ● ● ●
Tuber Blight	70	● ● ● ● ●
Alternaria	76	● ● ● ● ●
Common scab	52	● ● ● ● ●
Powdery scab	47	● ● ● ● ●
Spraing	76	● ● ● ● ●
PVY	39	● ● ● ● ●
Yntn tuber tolerance	96	● ● ● ● ●

PCN Resistance

Ro1/4	8	● ● ● ● ●
Ro2/3	*1	● ● ● ● ●
Ro5	*1	● ● ● ● ●
Pa2	*1	● ● ● ● ●
Pa3	*1	● ● ● ● ●

Wart disease

F1	10	● ● ● ● ●
F2	6	● ● ● ● ●
F6	6	● ● ● ● ●
F18		● ● ● ● ●

Fertilizer

- Adapt fertilization to soil analysis.
- Determinacy Group 3 for RB209
- Apply 2/3 before planting and 1/3 as top dressing.
- First top dressing of nitrogen directly after tuber initiation.
- Nitrogen (N) : High inputs (210-250Kg N/Ha inclusive of soil supply).
- Phosphate (P) and Potassium (K) : according to standard advice.

* HZPC own analysis/no official analysis



Production advice ware potatoes FRENCH FRIES



Pre-treatment and planting

- VICTORIA has a good dormancy.
- Allow the seed to acclimatize to the local conditions before planting.
- Always treat VICTORIA seeds and soil against Rhizoctonia.
- VICTORIA can be grown on all soil types.
- Plant seed free from condensation to prevent disease contamination.
- Planting with small white sprouts produces the best results.




Growing attention points

- VICTORIA grows spread in the ridge.
- Emergence is normal.
- VICTORIA is susceptible to metribuzin, also when applied before emergence.
- VICTORIA is high resistant to TRV.



Haulm killing and harvest

- The skin should be fully set before harvest.
- VICTORIA is moderately susceptible to bruising. Reduce drop heights and avoid mechanical damage.



Storage

- VICTORIA has a long dormancy and has good storability.
- Take care of the wound healing period.
- Store at a stable 8 °C.
- Ventilate regularly, but briefly, to prevent CO2 accumulation.
- For long term storage it is necessary to use sprout inhibitors.

- **Good storability**

