




## Production advice ware potatoes TRADITIONAL FRESH

- Big size tubers
- Very attractive bright skin
- Good yield
- Multi-purpose variety
- Good resistance package



### Agronomic characters

Maturity	59	Medium late
Yield early	106	High
Dormancy	70	Long
Yield mature	107	High
Tuber size	83	Large
Tuber shape		Long oval
Number of tubers		9-11
Skin colour		Light yellow
Flesh after cooking		Yellow
Cooking type		B - Slightly mealy
Dry matter content/Starch		20,9% / 15,1%
UWW/Specific gravity		385 / 1,083
Internal bruising	7	Little sensitive
Metribuzin sensitivity	83	● ● ● ● ●
Little Potato disorder	75	● ● ● ● ●



Skin and flesh colour



Cooking type



Maturity

### Plant populations

Seed size	Plant population/ha	Row distance	
		75 cm	90 cm
28/35	47.500	28	23
35/45	42.000	32	26
35/55	37.000	36	30
50/55	35.000	38	32
55/65	33.500	40	33

Plant depth: normal.

Always check the tuber count for an accurate calculation.

### Resistances

Foliage Blight	52	● ● ● ● ●
Tuber Blight	71	● ● ● ● ●
Alternaria	76	● ● ● ● ●
Common scab	57	● ● ● ● ●
Powdery scab	50	● ● ● ● ●
Spraing	55	● ● ● ● ●
PVY	12	● ● ● ● ●
Yntn tuber tolerance	99	● ● ● ● ●

#### PCN Resistance

Ro1/4 *1	● ● ● ● ●
Ro2/3 *1	● ● ● ● ●
Ro5 *1	● ● ● ● ●
Pa2 4	● ● ● ● ●
Pa3 3	● ● ● ● ●

#### Wart disease

F1 3	● ● ● ● ●
F2 *1	● ● ● ● ●
F6 4	● ● ● ● ●
F18 3	● ● ● ● ●

\* HZPC own analysis/no official analysis

### Fertilizer

- Adapt fertilization to soil analysis.
- Requires high nitrogen rates during early development.
- 100% of the nitrogen should be given at planting in one dose. This will help advance the crop.
- Nitrogen (N) : Medium inputs (170-210Kg N/Ha inclusive of soil supply).
- Phosphate (P) and Potassium (K) : according to standard advice.
- Don't use organic manure.



## Production advice ware potatoes TRADITIONAL FRESH



### Pre-treatment and planting

- VOYAGER has a long dormancy.
- A heat treatment is preferred to break the dormancy.
- Best grown on light, not too rich or even poor soils.
- Can be planted with small white buds but good pre-sprouting can advance the crop.
- Early planting is advised.




### Growing attention points

- The tuberisation is late.
- VOYAGER has a slow emergence but foliage develops strongly at a later stage.
- Use products that have an effect on *Alternaria solani* and *Alternaria alternata*.
- Despite the good *Phytophthora* resistance, a regular preventive application is advised given the long growing season.
- VOYAGER has a low susceptibility to common scab.
- VOYAGER is high resistant to TRV.



### Haulm killing and harvest

- Ensure the crop is naturally mature before haulm killing.



### Storage

- VOYAGER is little susceptible to *Fusarium*.
- Ventilate regularly, but briefly, to prevent CO<sub>2</sub> accumulation.
- Good to storable at constant temperature.
- VOYAGER has a long dormancy. Sprout inhibitors are only needed for late deliveries.